

# JEE Advanced Exam 2019 (Paper & Solution)

Date : 27 / 05 / 2019

## PAPER-1


### PART-I (CHEMISTRY)

#### SECTION – 1 (Maximum Marks : 12)


- This section contains **FOUR (04)** questions
- Each question has **FOUR** options. **ONLY ONE** of these four options is correct answer.
- For each question, choose the correct option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme :
  - Full Marks : +3 If **ONLY** the correct option is chosen.
  - Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered).
  - Negative Marks : -1 In all other cases.

**Q.1** Molar conductivity ( $\Lambda_m$ ) of aqueous solution of sodium stearate, which behaves as a strong electrolyte, is recorded at varying concentrations ( $c$ ) of sodium stearate. Which one of the following plots provides the correct representation of micelle formation in the solution ?

(critical micelle concentration (CMC) is marked with an arrow in the figures)



**Ans.** [3]



After CMC, as the conc. increases the aggregation of sodium stearate occurs &  $\Lambda_m$  decreases.


**Q.2** Calamine, malachite, magnetite and cryolite, respectively, are -

- (1)  $\text{ZnCO}_3$ ,  $\text{CuCO}_3$ ,  $\text{Fe}_2\text{O}_3$ ,  $\text{Na}_3\text{AlF}_6$                       (2)  $\text{ZnSO}_4$ ,  $\text{CuCO}_3$ ,  $\text{Fe}_2\text{O}_3$ ,  $\text{AlF}_3$   
 (3)  $\text{ZnSO}_4$ ,  $\text{Cu}(\text{OH})_2$ ,  $\text{Fe}_3\text{O}_4$ ,  $\text{Na}_3\text{AlF}_6$                       (4)  $\text{ZnCO}_3$ ,  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$ ,  $\text{Fe}_3\text{O}_4$ ,  $\text{Na}_3\text{AlF}_6$

**Ans.** [4]

**Sol.** Calamine =  $\text{ZnCO}_3$   
 Malachite =  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$   
 Magnetite =  $\text{Fe}_3\text{O}_4$   
 Cryolite =  $\text{Na}_3\text{AlF}_6$


**Q.3** The correct order of acid strength of the following carboxylic acids is -



- (1) I > II > III > IV      (2) III > II > I > IV      (3) II > I > IV > III      (4) I > III > II > IV

**Ans.** [1]

**Sol.**



I > II > III > IV

Acidic strength  $\propto$  stability of conjugate anion

**Q.4** The green colour produced in the borax bead test of a chromium (III) salt is due to

- (1)  $\text{Cr}(\text{BO}_2)_3$                       (2)  $\text{CrB}$                       (3)  $\text{Cr}_2\text{O}_3$                       (4)  $\text{Cr}_2(\text{B}_4\text{O}_7)_3$

**Ans.** [1]

**Sol.**  $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O} \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7 \xrightarrow{\Delta} \text{NaBO}_2 + \text{B}_2\text{O}_3$   
 $\text{Cr}^{3+} + \text{B}_2\text{O}_3 \rightarrow \text{Cr}(\text{BO}_2)_3$   
 Green

**SECTION – 2 (Maximum Marks : 32)**




- This section contains **EIGHT (08)** questions
- Each question has **FOUR** options. **ONE OR MORE THAN ONE** of these four option(s) is (are) correct option(s).
- For each question, choose(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme :
 

Full Marks	: +4	If only (all) the correct option(s) is (are) chosen.
Partial Marks	: +3	If all the four options are correct but <b>ONLY</b> three options are chosen.
Partial Marks	: +2	If three or more options are correct but <b>ONLY</b> two options are chosen, both of which are correct options.
Partial Marks	: +1	If two or more options are correct but <b>ONLY</b> one option is chosen and it is a correct option.
Zero Marks	: 0	If none of the option is chosen (i.e. the question is unanswered).
Negative Marks	: -1	In all other cases.





**Q.1** Which of the following statement(s) is (are) true ?

- (1) The two six-membered cyclic hemiacetal forms of D-(+)- glucose are called anomers
- (2) Hydrolysis of sucrose gives dextrorotatory glucose and laevorotatory fructose
- (3) Monosaccharides cannot be hydrolysed to give polyhydroxy aldehydes and ketones
- (4) Oxidation of glucose with bromine water gives glutamic acid


**Ans.** [1, 2, 3]




**Q.2** Choose the reaction(s) from the following option, for which the standard enthalpy of reaction is equal to the standard enthalpy of formation-



**Ans.** [1,4]




**Q.3** An tin chloride Q undergoes the following reactions (not balanced)



X is a monanion having pyramidal geometry. Both Y and Z are neutral compounds. Choose the correct options (s)




- (1) The oxidation state of the central atom in Z is +2
- (2) There is a coordinate bond in Y
- (3) The central atom in Z has one lone pair of electrons
- (4) The central atom in X is sp<sup>3</sup> hybridized

**Ans.** [2, 4]




**Q.4** Fusion of MnO<sub>2</sub> with KOH in presence of O<sub>2</sub> produces a salt W. Alkaline solution of W upon electrolytic oxidation yields another salt X. The manganese containing ions present in W and X, respectively are Y and Z. Correct statement (s) is (are) -

- (1) Both Y and Z are coloured and have tetrahedral shape
- (2) Y is diamagnetic in nature while Z is paramagnetic
- (3) In both Y and Z, π-bonding occurs between p-orbitals of oxygen and d-orbitals of manganese
- (4) In aqueous acidic solution, Y undergoes disproportionation reaction to give Z and MnO<sub>2</sub>

**Ans. [1,3,4]**


$\text{MnO}_4^-$  (Z) = +7 oxidation state = 0 unpaired  $e^-$  diamagnetic  
+7

$\text{MnO}_4^{2-}$  (Y) = +6 oxidation state = 1 unpaired  $e^-$  paramagnetic



**Q.5** Which of the following statement(s) is (are) correct regarding the root mean square speed ( $U_{\text{rms}}$ ) and average translational kinetic energy ( $\epsilon_{\text{av}}$ ) of a molecule in a gas at equilibrium ?

- (1)  $\epsilon_{\text{av}}$  is doubled when its temperature is increased four times
- (2)  $\epsilon_{\text{av}}$  at a given temperature does not depend on its molecular mass
- (3)  $U_{\text{rms}}$  is doubled when its temperature is increased four times
- (4)  $U_{\text{rms}}$  is inversely proportional to the square root of its molecular mass


**Ans. [2, 3, 4]**



**Sol.**  $E_{\text{av}} = \frac{3}{2}KT$

$U_{\text{rms}} = \sqrt{\frac{3RT}{m}}$


**Q.6** Each of the following option contains a set of four molecules, Identify the option(s) where all four molecules possess permanent dipole moment at room temperature -

- (1)  $\text{SO}_2$ ,  $\text{C}_6\text{H}_5\text{Cl}$ ,  $\text{H}_2\text{Se}$ ,  $\text{BrF}_5$
- (2)  $\text{BeCl}_2$ ,  $\text{CO}_2$ ,  $\text{BCl}_3$ ,  $\text{CHCl}_3$
- (3)  $\text{BF}_3$ ,  $\text{O}_3$ ,  $\text{SF}_6$ ,  $\text{XeF}_6$
- (4)  $\text{NO}_2$ ,  $\text{NH}_3$ ,  $\text{POCl}_3$ ,  $\text{CH}_3\text{Cl}$

**Ans. [1, 4]**







**Q.7** In the decay sequence



$x_1, x_2, x_3$  and  $x_4$  are particles / radiation emitted by the respective isotopes. The correct option(s) is(are)

- (1)  $x_3$  is  $\gamma$ -ray      (2)  $x_2$  is  $\beta^-$   
 (3) Z is an isotope of uranium      (4)  $x_1$  will deflect towards negatively charged plate

**Ans.** [2, 3, 4]  
**Sol.**














$$x_1 = \alpha$$

$$x_2 = \beta$$


$$x_3 = \beta$$

$$x_4 = \alpha$$

**Q.8** Choose the correct option(s) for the following set of reactions



**Ans.** []

**Sol.**


### SECTION – 3 (Maximum Marks : 18)

- This section contains **SIX (06)** questions. The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme.  
 Full Marks : +3 If **ONLY** the correct numerical value is entered as answer.  
 Zero Marks : 0 In all other cases.

**Q.1** Consider the kinetic data given in the following table for the reaction  $A + B + C \rightarrow \text{Product}$

Experiment No.	[A] (mol dm <sup>-3</sup> )	[B] (mol dm <sup>-3</sup> )	[C] (mol dm <sup>-3</sup> )	Rate of reaction (mol dm <sup>-3</sup> s <sup>-1</sup> )
1	0.2	0.1	0.1	$6.0 \times 10^{-5}$
2	0.2	0.2	0.1	$6.0 \times 10^{-5}$
3	0.2	0.1	0.2	$1.2 \times 10^{-4}$
4	0.3	0.1	0.1	$9.0 \times 10^{-5}$

The rate of the reaction for  $[A] = 0.15 \text{ mol dm}^{-3}$ ,  $[B] = 0.25 \text{ mol dm}^{-3}$  and  $[C] = 0.15 \text{ mol dm}^{-3}$  is found to be  $Y \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$ . The value of Y is \_\_\_\_\_ .

**Ans. [6.75]**
**Sol.**  $r = K[A]^p[B]^q[C]^n$ 

$$\frac{r_2}{r_1} = \left[ \frac{B_2}{B_1} \right]^q$$

$$1 = 2^q \quad \text{or} \quad 2^0 = 2^q \quad \therefore q = 0$$

$$\frac{r_3}{r_2} = \left[ \frac{C_3}{C_2} \right]^n$$

$$2 = 2^n \quad n = 1$$

$$\frac{r_4}{r_1} = \left[ \frac{A_4}{A_1} \right]^p \left[ \frac{C_4}{C_1} \right]^n$$

$$\frac{9}{6} = \left(\frac{3}{2}\right)^P \times \left(\frac{3}{2}\right)^1 \Rightarrow P = 1$$


$$r = K[A]^1[C]^1$$

$$K = \frac{r}{[A][C]} = \frac{6 \times 10^{-5}}{2 \times 10^{-2}} = 3 \times 10^{-3}$$

$$\begin{aligned} r &= K[A][C] \\ &= 3 \times 10^{-3} \times 0.15 \times 0.15 \\ &= 6.75 \times 10^{-5} \end{aligned}$$



**Q.2** Among  $B_2H_6$ ,  $B_3N_3H_6$ ,  $N_2O$ ,  $N_2O_4$ ,  $H_2S_2O_3$  and  $H_2S_2O_8$  the total number of molecules containing covalent bond between two atoms of the same kind is \_\_\_\_\_

**Ans. [4]**



**Q.3** At 143 K, the reaction of  $XeF_4$  with  $O_2F_2$  produces a xenon compound Y. The total number of lone pair(s) of electron present on the whole molecule of Y is \_\_\_\_\_

**Ans. [19]**



Total no. of lone pair present on the whole molecule of Y = 1 l. p. in Xe + 18. l. p. in F  
= 19 total l. p.





**Q.4** On dissolving 0.5 g of a non-volatile non-ionic solute to 39 g of benzene, its vapor pressure decreases from 650 mm Hg to 640 mm Hg. The depression of freezing point of benzene (in K) upon addition of the solute is \_\_\_\_\_

(Given data : Molar mass and the molal freezing point depression constant of benzene are 78 g mol<sup>-1</sup> and 5.12 K kg mol<sup>-1</sup>, respectively)

**Ans. [1.03]**

**Sol.**

$$\frac{P_B^0 - P_S}{P_S} = \frac{n_A}{n_B}$$

$$\frac{650 - 640}{640} = \frac{n_A}{0.5}$$

$$\frac{10 \times 0.5}{640} = n_A = \frac{5}{640}$$


$$\Delta T_f = iK_f m$$

$$= (1)(5.12) \frac{5 \times 1000}{640 \times 39}$$

$$= 1.0256 = 1.026$$


$$= 1.03$$

**Q.5** For the following reaction the equilibrium constant K<sub>c</sub> at 298 K is 1.6 × 10<sup>17</sup>



When equal volumes of 0.06 M Fe<sup>2+</sup> (aq) and 0.2 M S<sup>2-</sup> (aq) solutions are mixed, the equilibrium concentration of Fe<sup>2+</sup> (aq) is found to be Y × 10<sup>-17</sup> M. The value of Y is \_\_\_\_\_

**Ans. [8.93]**



Since,

$$K_c = K_c = 1.6 \times 10^{17} \text{ (Very high value)}$$

$$K_c = \frac{1}{[\text{Fe}^{+2}][\text{S}^{-2}]}$$

$$\delta = [\text{Fe}^{+2}] = \frac{1}{[K_c][\text{S}^{-2}]}$$

$$= \frac{1}{(1.6 \times 10^{17})(0.07)}$$


$$= \frac{100}{7 \times 1.6} \times 10^{-17}$$

$$= 8.93 \times 10^{-17}$$


$$Y = 8.93$$

**Q.6** Schemes 1 and 2 describe the conversion of P to Q and R to S, respectively. Scheme 3 describes the synthesis of T from Q and S. The total number of Br atoms in a molecule of T is \_\_\_\_\_


**Scheme 1 :**



**Scheme 2 :**




**Scheme 3 :**




**Ans. [4]**

**Sol. Scheme 1 :**



**Scheme 2 :**



**Scheme 3 :**

