

**TEST PAPER OF JEE(MAIN) EXAMINATION – 2019**  
**(Held On Thursday 10<sup>th</sup> JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM**  
**CHEMISTRY**

1. An ideal gas undergoes isothermal compression from  $5 \text{ m}^3$  against a constant external pressure of  $4 \text{ Nm}^{-2}$ . Heat released in this process is used to increase the temperature of 1 mole of Al. If molar heat capacity of Al is  $24 \text{ J mol}^{-1} \text{ K}^{-1}$ , the temperature of Al increases by :

- (1)  $\frac{3}{2} \text{ K}$     (2)  $\frac{2}{3} \text{ K}$     (3) 1 K    (4) 2 K

Ans. (2)

2. The 71<sup>st</sup> electron of an element X with an atomic number of 71 enters into the orbital :

- (1) 4f    (2) 6p    (3) 6s    (4) 5d

Ans. (1)

3. The number of 2-centre-2-electron and 3-centre-2-electron bonds in  $\text{B}_2\text{H}_6$ , respectively, are :

- (1) 2 and 4                      (2) 2 and 1  
 (3) 2 and 2                      (4) 4 and 2

Ans. (4)

4. The amount of sugar ( $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ ) required to prepare 2 L of its 0.1 M aqueous solution is :

- (1) 68.4 g    (2) 17.1 g    (3) 34.2 g    (4) 136.8 g

Ans. (1)

5. Among the following reactions of hydrogen with halogens, the one that requires a catalyst is :

- (1)  $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$     (2)  $\text{H}_2 + \text{F}_2 \rightarrow 2\text{HF}$   
 (3)  $\text{H}_2 + \text{Cl}_2 \rightarrow 2\text{HCl}$     (4)  $\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$






Ans. (1)

6. Sodium metal on dissolution in liquid ammonia gives a deep blue solution due to the formation of:

- (1) sodium ion-ammonia complex  
 (2) sodamide  
 (3) sodium-ammonia complex  
 (4) ammoniated electrons

Ans. (4)

7. What will be the major product in the following mononitration reaction ?



Ans. (3)




8. In the cell  $\text{Pt(s)}|\text{H}_2(\text{g}, 1\text{bar})|\text{HCl}(\text{aq})|\text{Ag(s)}|\text{Pt(s)}$  the cell potential is 0.92 when a  $10^{-6}$  molal HCl solution is used. The standard electrode potential of  $(\text{AgCl}/\text{Ag}, \text{Cl}^-)$  electrode is :

$$\left\{ \text{given, } \frac{2.303RT}{F} = 0.06 \text{ V at } 298 \text{ K} \right\}$$

- (1) 0.20 V    (2) 0.76 V    (3) 0.40 V    (4) 0.94 V

Ans. (1)

9. The major product of the following reaction is:








Ans. (3)

10. The pair that contains two P-H bonds in each of the oxoacids is :

- (1)  $\text{H}_3\text{PO}_2$  and  $\text{H}_4\text{P}_2\text{O}_5$
- (2)  $\text{H}_4\text{P}_2\text{O}_5$  and  $\text{H}_4\text{P}_2\text{O}_6$
- (3)  $\text{H}_3\text{PO}_3$  and  $\text{H}_3\text{PO}_2$
- (4)  $\text{H}_4\text{P}_2\text{O}_5$  and  $\text{H}_3\text{PO}_3$

Ans. (1)

11. The major product of the following reaction is:



Ans. (4)

12. The difference in the number of unpaired electrons of a metal ion in its high-spin and low-spin octahedral complexes is two. The metal ion is :

- (1)  $\text{Fe}^{2+}$
- (2)  $\text{Co}^{2+}$
- (3)  $\text{Mn}^{2+}$
- (4)  $\text{Ni}^{2+}$

Ans. (2)

13. A compound of formula  $\text{A}_2\text{B}_3$  has the hcp lattice. Which atom forms the hcp lattice and what fraction of tetrahedral voids is occupied by the other atoms :

- (1) hcp lattice-A,  $\frac{2}{3}$  Tetrahedral voids-B





- (2) hcp lattice-B,  $\frac{1}{3}$  Tetrahedral voids-A

- (3) hcp lattice-B,  $\frac{2}{3}$  Tetrahedral voids-A

- (4) hcp lattice-A,  $\frac{1}{3}$  Tetrahedral voids-B

Ans. (2)

14. The reaction that is NOT involved in the ozone layer depletion mechanism in the stratosphere is:



Ans. (3)

15. The process with negative entropy change is :

- (1) Dissolution of iodine in water
- (2) Synthesis of ammonia from  $\text{N}_2$  and  $\text{H}_2$
- (3) Dissolution of  $\text{CaSO}_4(\text{s})$  to  $\text{CaO}(\text{s})$  and  $\text{SO}_3(\text{g})$
- (4) Sublimation of dry ice

Ans. (2)

**MAJOR COMPUTER BASED TEST (CBT) SERIES**


**JEE (Advanced)- Target 2019**





[dlp.allen.ac.in](http://dlp.allen.ac.in)

Test Dates: 3<sup>rd</sup> Feb, 21<sup>st</sup> & 28<sup>th</sup> April, 12<sup>th</sup> May

0744-2750275

16. The major product of the following reaction is:



- (1) 
- (2) 
- (3) 
- (4) 


Ans. (3)





17. A reaction of cobalt(III) chloride and ethylenediamine in a 1 : 2 mole ratio generates two isomeric products A (violet coloured) B (green coloured). A can show optical activity, B is optically inactive. What type of isomers does A and B represent ?

- (1) Geometrical isomers  
 (2) Ionisation isomers  
 (3) Coordination isomers  
 (4) Linkage isomers

Ans. (1)

18. The major product obtained in the following reaction is :



- (1) 
- (2) 
- (3) 
- (4) 


Ans. (4)

19. Which of the following tests cannot be used for identifying amino acids ?

- (1) Biuret test                      (2) Xanthoproteic test  
 (3) Barfoed test                      (4) Ninhydrin test

Ans. (3)

20. What is the IUPAC name of the following compound ?



- (1) 3-Bromo-1, 2-dimethylbut-1-ene]  
 (2) 4-Bromo-3-methylpent-2-ene  
 (3) 2-Bromo-3-methylpent-3-ene  
 (4) 3-Bromo-3-methyl-1, 2-dimethylprop-1-ene

Ans. (2)

**MAJOR COMPUTER BASED TEST (CBT) SERIES**


**JEE (Main)- Target 2019**

dlp.allen.ac.in

Test Dates: 24<sup>th</sup> & 31<sup>st</sup> March

0744-2750275

21. Which is the most suitable reagent for the following transformation ?



- (1) alkaline  $\text{KMnO}_4$       (2)  $\text{I}_2/\text{NaOH}$   
(3) Tollen's reagent      (4)  $\text{CrO}_2/\text{CS}_2$

Ans. (2)

22. The correct match between item 'I' and item 'II' is :

Item 'I' (compound)	Item 'II' (reagent)
(A) Lysine	(P) 1-naphthol
(B) Furfural	(Q) ninhydrin
(C) Benzyl alcohol	(R) $\text{KMnO}_4$
(D) Styrene	(S) Ceric ammonium nitrate

- (1) (A)→(Q), (B)→(P), (C)→(S), (D)→(R)  
(2) (A)→(Q), (B)→(R), (C)→(S), (D)→(P)  
(3) (A)→(Q), (B)→(P), (C)→(R), (D)→(S)  
(4) (A)→(R), (B)→(P), (C)→(Q), (D)→(S)

Ans. (1)

23. In the reaction of oxalate with permanganate in acidic medium, the number of electrons involved in producing one molecule of  $\text{CO}_2$  is :

- (1) 10      (2) 2      (3) 1      (4) 5

Ans. (3)

24. 5.1g  $\text{NH}_4\text{SH}$  is introduced in 3.0 L evacuated flask at  $327^\circ\text{C}$ . 30% of the solid  $\text{NH}_4\text{SH}$  decomposed to  $\text{NH}_3$  and  $\text{H}_2\text{S}$  as gases. The  $K_p$  of the reaction at  $327^\circ\text{C}$  is ( $R = 0.082 \text{ L atm mol}^{-1}\text{K}^{-1}$ , Molar mass of S =  $32 \text{ g mol}^{-1}$ , molar mass of N =  $14 \text{ g mol}^{-1}$ )

- (1)  $1 \times 10^{-4} \text{ atm}^2$       (2)  $4.9 \times 10^{-3} \text{ atm}^2$   
(3)  $0.242 \text{ atm}^2$       (4)  $0.242 \times 10^{-4} \text{ atm}^2$

Ans. (3)

25. The electrolytes usually used in the electroplating of gold and silver, respectively, are :

- (1)  $[\text{Au}(\text{OH})_4]^-$  and  $[\text{Ag}(\text{OH})_2]^-$   
(2)  $[\text{Au}(\text{CN})_2]^-$  and  $[\text{Ag} \text{Cl}_2]^-$   
(3)  $[\text{Au}(\text{NH}_3)_2]^+$  and  $[\text{Ag}(\text{CN})_2]^-$   
(4)  $[\text{Au}(\text{CN})_2]^-$  and  $[\text{Ag}(\text{CN})_2]^-$





Ans. (4)

26. Elevation in the boiling point for 1 molal solution of glucose is 2 K. The depression in the freezing point of 2 molal solutions of glucose in the same solvent is 2 K. The relation between  $K_b$  and  $K_f$  is:

- (1)  $K_b = 0.5 K_f$       (2)  $K_b = 2 K_f$   
(3)  $K_b = 1.5 K_f$       (4)  $K_b = K_f$

Ans. (2)

27. An aromatic compound 'A' having molecular formula  $\text{C}_7\text{H}_6\text{O}_2$  on treating with aqueous ammonia and heating forms compound 'B'. The compound 'B' on reaction with molecular bromine and potassium hydroxide provides compound 'C' having molecular formula  $\text{C}_6\text{H}_7\text{N}$ . The structure of 'A' is :




Ans. (3)

28. The ground state energy of hydrogen atom is  $-13.6 \text{ eV}$ . The energy of second excited state  $\text{He}^+$  ion in eV is :

- (1)  $-6.04$       (2)  $-27.2$       (3)  $-54.4$       (4)  $-3.4$

Ans. (1)

29. For an elementary chemical reaction,



- (1)  $2k_1[\text{A}_2]-k_{-1}[\text{A}]^2$       (2)  $k_1[\text{A}_2]-k_{-1}[\text{A}]^2$   
(3)  $2k_1[\text{A}_2]-2k_{-1}[\text{A}]^2$       (4)  $k_1[\text{A}_2]+k_{-1}[\text{A}]^2$

Ans. (3)

30. Haemoglobin and gold sol are examples of :

- (1) negatively charged sols  
(2) positively charged sols  
(3) negatively and positively charged sols, respectively  
(4) positively and negatively charged sols, respectively

Ans. (4)