

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019

(Held On Friday 11<sup>th</sup> JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM

MATHEMATICS

1. Let  $A = \begin{pmatrix} 0 & 2q & r \\ p & q & -r \\ p & -q & r \end{pmatrix}$ . If  $AA^T = I_3$ , then  $|p|$

is :

(1)  $\frac{1}{\sqrt{2}}$  (2)  $\frac{1}{\sqrt{5}}$

(3)  $\frac{1}{\sqrt{6}}$  (4)  $\frac{1}{\sqrt{3}}$

Ans. (1)

2. The area (in sq. units) of the region bounded by the curve  $x^2 = 4y$  and the straight line  $x = 4y - 2$  :-

(1)  $\frac{5}{4}$  (2)  $\frac{9}{8}$

(3)  $\frac{3}{4}$  (4)  $\frac{7}{8}$

Ans. (2)

3. The outcome of each of 30 items was observed;

10 items gave an outcome  $\frac{1}{2} - d$  each, 10 items

gave outcome  $\frac{1}{2}$  each and the remaining

10 items gave outcome  $\frac{1}{2} + d$  each. If the

variance of this outcome data is  $\frac{4}{3}$  then  $|d|$  equals :-

(1) 2 (2)  $\frac{\sqrt{5}}{2}$

(3)  $\frac{2}{3}$  (4)  $\sqrt{2}$

Ans. (4)

4. The sum of an infinite geometric series with positive terms is 3 and the sum of the cubes of its terms is  $\frac{27}{19}$ . Then the common ratio of this series is :

(1)  $\frac{4}{9}$  (2)  $\frac{2}{9}$

(3)  $\frac{2}{3}$  (4)  $\frac{1}{3}$

Ans. (3)

5. Let  $\vec{a} = \hat{i} + 2\hat{j} + 4\hat{k}$ ,  $\vec{b} = \hat{i} + \lambda\hat{j} + 4\hat{k}$  and  $\vec{c} = 2\hat{i} + 4\hat{j} + (\lambda^2 - 1)\hat{k}$  be coplanar vectors.

Then the non-zero vector  $\vec{a} \times \vec{c}$  is :

(1)  $-14\hat{i} - 5\hat{j}$  (2)  $-10\hat{i} - 5\hat{j}$

(3)  $-10\hat{i} + 5\hat{j}$  (4)  $-14\hat{i} + 5\hat{j}$

Ans. (3)

6. Let  $\left(-2 - \frac{1}{3}i\right)^3 = \frac{x + iy}{27}$  ( $i = \sqrt{-1}$ ), where  $x$

and  $y$  are real numbers, then  $y - x$  equals :

(1) -85 (2) 85

(3) -91 (4) 91

Ans. (4)

7. Let  $f(x) = \begin{cases} -1, & -2 \leq x < 0 \\ x^2 - 1, & 0 \leq x \leq 2 \end{cases}$  and

$g(x) = |f(x)| + f(|x|)$ . Then, in the interval  $(-2, 2)$ ,  $g$  is :-

(1) differentiable at all points

(2) not differentiable at two points

(3) not continuous

(4) not differentiable at one point

Ans. (4)

8. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $f(x) = \frac{x}{1+x^2}$ ,

$x \in \mathbb{R}$ . Then the range of  $f$  is :

- (1)  $(-1, 1) - \{0\}$       (2)  $\left[-\frac{1}{2}, \frac{1}{2}\right]$   
 (3)  $\mathbb{R} - \left[-\frac{1}{2}, \frac{1}{2}\right]$       (4)  $\mathbb{R} - [-1, 1]$

Ans. (2)

9. The sum of the real values of  $x$  for which the middle term in the binomial expansion of

$\left(\frac{x^3}{3} + \frac{3}{x}\right)^8$  equals 5670 is :

- (1) 6      (2) 8      (3) 0      (4) 4

Ans. (3)

10. The value of  $r$  for which

${}^{20}C_r + {}^{20}C_0 + {}^{20}C_{r-1} + {}^{20}C_1 + {}^{20}C_{r-2} + {}^{20}C_2 + \dots + {}^{20}C_0$  is maximum, is

- (1) 20      (2) 15  
 (3) 11      (4) 10

Ans. (1)

11. Let  $a_1, a_2, \dots, a_{10}$  be a G.P. If  $\frac{a_3}{a_1} = 25$ , then

$\frac{a_9}{a_5}$  equals :

- (1)  $2(5^2)$       (2)  $4(5^2)$   
 (3)  $5^4$       (4)  $5^3$

Ans. (3)

12. If  $\int \frac{\sqrt{1-x^2}}{x^4} dx = A(x) \left(\sqrt{1-x^2}\right)^m + C$ , for

a suitable chosen integer  $m$  and a function  $A(x)$ , where  $C$  is a constant of integration then  $(A(x))^m$  equals :

- (1)  $\frac{-1}{3x^3}$       (2)  $\frac{-1}{27x^9}$   
 (3)  $\frac{1}{9x^4}$       (4)  $\frac{1}{27x^6}$

Ans. (2)

13. In a triangle, the sum of lengths of two sides is  $x$  and the product of the lengths of the same two sides is  $y$ . If  $x^2 - c^2 = y$ , where  $c$  is the length of the third side of the triangle, then the circumradius of the triangle is :

- (1)  $\frac{y}{\sqrt{3}}$       (2)  $\frac{c}{\sqrt{3}}$       (3)  $\frac{c}{3}$       (4)  $\frac{3}{2}y$

Ans. (2)

14. The value of the integral  $\int_{-2}^2 \frac{\sin^2 x}{\left[\frac{x}{\pi}\right] + \frac{1}{2}} dx$

(where  $[x]$  denotes the greatest integer less than  ${}^{20}C_r$  or equal to  $x$ ) is :

- (1) 4      (2)  $4 - \sin 4$   
 (3)  $\sin 4$       (4) 0

Ans. (4)

15. If the system of linear equations

$$\begin{aligned} 2x + 2y + 3z &= a \\ 3x - y + 5z &= b \\ x - 3y + 2z &= c \end{aligned}$$

where  $a, b, c$  are non-zero real numbers, has more than one solution, then :

- (1)  $b - c - a = 0$       (2)  $a + b + c = 0$   
 (3)  $b + c - a = 0$       (4)  $b - c + a = 0$

Ans. (1)

16. A square is inscribed in the circle

$x^2 + y^2 - 6x + 8y - 103 = 0$  with its sides parallel to the coordinate axes. Then the distance of the vertex of this square which is nearest to the origin is :-

- (1) 13      (2)  $\sqrt{137}$   
 (3) 6      (4)  $\sqrt{41}$

Ans. (4)

17. Let  $f_k(x) = \frac{1}{k}(\sin^k x + \cos^k x)$  for  $k = 1, 2,$

$3, \dots$ . Then for all  $x \in \mathbb{R}$ , the value of  $f_4(x) - f_6(x)$  is equal to :-

- (1)  $\frac{5}{12}$       (2)  $\frac{-1}{12}$       (3)  $\frac{1}{4}$       (4)  $\frac{1}{12}$

Ans. (4)

18. Let  $[x]$  denote the greatest integer less than or equal to  $x$ . Then :-

$$\lim_{x \rightarrow 0} \frac{\tan(\pi \sin^2 x) + (|x| - \sin(x[x]))^2}{x^2}$$

- (1) equals  $\pi$                       (2) equals 0  
 (3) equals  $\pi + 1$                 (4) does not exist

**Ans. (4)**

19. The direction ratios of normal to the plane through the points  $(0, -1, 0)$  and  $(0, 0, 1)$  and

making an angle  $\frac{\pi}{4}$  with the plane  $y-z+5=0$  are:

- (1)  $2\sqrt{3}, 1, -1$                 (2)  $2, \sqrt{2}, -\sqrt{2}$   
 (3)  $2, -1, 1$                       (4)  $\sqrt{2}, 1, -1$

**Ans. (2, 4)**

20. If  $x \log_e(\log_e x) - x^2 + y^2 = 4 (y > 0)$ , then  $dy/dx$  at  $x = e$  is equal to :

(1)  $\frac{e}{\sqrt{4+e^2}}$                       (2)  $\frac{(1+2e)}{2\sqrt{4+e^2}}$

(3)  $\frac{(2e-1)}{2\sqrt{4+e^2}}$                       (4)  $\frac{(1+2e)}{\sqrt{4+e^2}}$

**Ans. (3)**

21. The straight line  $x + 2y = 1$  meets the coordinate axes at A and B. A circle is drawn through A, B and the origin. Then the sum of perpendicular distances from A and B on the tangent to the circle at the origin is :

(1)  $\frac{\sqrt{5}}{4}$     (2)  $\frac{\sqrt{5}}{2}$     (3)  $2\sqrt{5}$     (4)  $4\sqrt{5}$

**Ans. (2)**

22. If  $q$  is false and  $p \wedge q \leftrightarrow r$  is true, then which one of the following statements is a tautology?

- (1)  $(p \vee r) \rightarrow (p \wedge r)$   
 (2)  $p \vee r$   
 (3)  $p \wedge r$   
 (4)  $(p \wedge r) \rightarrow (p \vee r)$

**Ans. (4)**

23. If  $y(x)$  is the solution of the differential equation

$$\frac{dy}{dx} + \left( \frac{2x+1}{x} \right) y = e^{-2x}, \quad x > 0,$$

where  $y(1) = \frac{1}{2}e^{-2}$ , then :

(1)  $y(x)$  is decreasing in  $(0, 1)$

(2)  $y(x)$  is decreasing in  $\left(\frac{1}{2}, 1\right)$

(3)  $y(\log_e 2) = \frac{\log_e 2}{4}$

(4)  $y(\log_e 2) = \log_e 4$

**Ans. (2)**

24. The maximum value of the function

$f(x) = 3x^3 - 18x^2 + 27x - 40$  on the set

$S = \{x \in \mathbb{R} : x^2 + 30 \leq 11x\}$  is :

- (1) 122                                  (2) -222  
 (3) -122                                (4) 222

**Ans. (1)**

25. If one real root of the quadratic equation  $81x^2 + kx + 256 = 0$  is cube of the other root, then a value of  $k$  is

- (1) -81    (2) 100    (3) -300    (4) 144

**Ans. (3)**

26. Two circles with equal radii are intersecting at the points  $(0, 1)$  and  $(0, -1)$ . The tangent at the point  $(0, 1)$  to one of the circles passes through the centre of the other circle. Then the distance between the centres of these circles is :

- (1) 1                                      (2)  $\sqrt{2}$   
 (3)  $2\sqrt{2}$                               (4) 2

**Ans. (4)**

**MAJOR COMPUTER BASED TEST (CBT) SERIES**

**JEE (Advanced)- Target 2019**

**dlp.allen.ac.in**    Test Dates: 3<sup>rd</sup> Feb, 21<sup>st</sup> & 28<sup>th</sup> April, 12<sup>th</sup> May    **0744-2750275**

27. Equation of a common tangent to the parabola  $y^2 = 4x$  and the hyperbole  $xy = 2$  is :

- (1)  $x + 2y + 4 = 0$
- (2)  $x - 2y + 4 = 0$
- (3)  $x + y + 1 = 0$
- (4)  $4x + 2y + 1 = 0$

Ans. (1)

28. The plane containing the line  $\frac{x-3}{2} = \frac{y+2}{-1} = \frac{z-1}{3}$

and also containing its projection on the plane  $2x + 3y - z = 5$ , contains which one of the following points ?

- (1) (2, 0, -2)
- (2) (-2, 2, 2)
- (3) (0, -2, 2)
- (4) (2, 2, 0)

Ans. (1)

29. If tangents are drawn to the ellipse  $x^2 + 2y^2 = 2$  at all points on the ellipse other than its four vertices then the mid points of the tangents intercepted between the coordinate axes lie on the curve :

- (1)  $\frac{x^2}{2} + \frac{y^2}{4} = 1$
- (2)  $\frac{x^2}{4} + \frac{y^2}{2} = 1$
- (3)  $\frac{1}{2x^2} + \frac{1}{4y^2} = 1$
- (4)  $\frac{1}{4x^2} + \frac{1}{2y^2} = 1$

Ans. (3)

30. Two integers are selected at random from the set  $\{1, 2, \dots, 11\}$ . Given that the sum of selected numbers is even, the conditional probability that both the numbers are even is :

- (1)  $\frac{2}{5}$
- (2)  $\frac{1}{2}$
- (3)  $\frac{3}{5}$
- (4)  $\frac{7}{10}$

Ans. (1)

**MAJOR COMPUTER BASED TEST (CBT) SERIES**

**JEE (Main)- Target 2019**

[dlp.allen.ac.in](http://dlp.allen.ac.in)

Test Dates: 24<sup>th</sup> & 31<sup>st</sup> March

0744-2750275